

Erectile dysfunction therapy after nerve sparing prostatectomy: six month comparison between avanafil 200 mg and sildenafil 100 mg

Pier Andrea Della Camera 1, Mauro Gacci 1, Vincenzo Maria Altieri 2, Emanuele Micheli 2, Giovanni Tasso 1, Giulia Bencini 1, Gianmartin Cito 1, Simone Morselli 1, Nicola Laruccia 1, Andrea Cocci 1, Michele Marzocco 1, Fabrizio Travaglini 1, Sabino Scelzi 1, Antonio Rocco Mottola 3, Marco Carini

1 Azienda Ospedaliera Universitaria Careggi, Firenze, Italia 2 Humanitas Gavezzeni, Bergamo, Italia, 3 Azienda




	Sildenafil (n=80)		Avanafil (n=80)	
	Mean	±	Mean	±
Age	66yr	59-72 yr	64 yr	57-71 yr
Bmi	25,90	30,29-21,20	26,15	32,49-21,53
Waist circonference	103,26 cm	89-120 cm	98,61 cm	85-115 cm
Weight	79,92 kg	65-120 kg	81,9 kg	110-69 kg
Height	175,76 cm	180-164 cm	176,06 cm	198-170 cm
	Sildenafil (n=80)		Avanafil (n=80)	
	Number	%	Number	%
Hypertension	32	40%	24	30%
Diabetes	10	12,5%	13	16,25%
Hypertriglicerimia	13	16,25%	9	11,25%
Smokers	20	25%	12	15%
CCI-score 0	83	78,75%	80	75%
CCI-score 1 -Myocardial infarction=	17	21,25%	20	25%
-Diabetes (active therapy)=	8		10	
-Chronic Pulmonary disease=	8		9	
-Connective Tissue disease=	0		1	
Ecog-score 0	96	95%	96	95%
Ecog-score 1	4	5%	4	5%


Value at baseline	Sildenafil (n=80) mean	±	Avanafil (n=80) mean	±	P
IIEF 15	22,35	19-25	21,80	18-25	0,513
QoL	1,4	0-3	1,04	1-3	0,642
Weekly sexual intercourse	1,65	1-3	1,027	0,50-3	<0,001
Ehs	3,6	3-4	3,52	2-4	0,866

Preoperative setting


Epidemiological data





Postoperative IIEF




Δ Variation of IIEF between pre and postoperative




a) EHS b) Δ Variation preoperative/postoperative




a) Sexual quality of life. b) Δ variation of sexual Qol preoperative vs postoperative




a) Number of weekly sexual intercourse. b) Δ variation of number of sexual intercourse



a) Number of weekly sexual intercourse. b) Δ variation of number of sexual intercourse



**CONGRESSO
NAZIONALE
SIA 42° ANNO**




Conclusion:

Sep-Gaq Tests

-Avanafil 200 mg compared to the same therapy with Sildenafil 100 mg three times a week showed a lower ability to allow an effective erection in the initial phase of sexual intercourse and a lower penile hardness

-Avanafil 200mg showed greater safety in its administration showing a lower presence of side effects compared to Sildenafil 100 mg



ADVERSE EVENTS			
	%	Discontinued	%
Sildenafil	16	9	11,25%
Avanafil	4	0	0%
Headache	8	4	5%
Fainting	6	0	1,25%
Dyspepsia	3	0	0%
Cyanopsia	1	0	0%

Thanks for the attention

